

Long-Term Ambient Air Pollution and Childhood Eczema in the United States

Joshua P. Keller,¹ Joan H. Dunlop,² Nathan A. Ryder,¹ Roger D. Peng,³ and Corinne A. Keet⁴

¹Department of Statistics, Colorado State University, Fort Collins, Colorado, USA

²Division of Pediatric Allergy and Immunology, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

³Department of Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

⁴Division of Pediatric Allergy and Immunology, University of North Carolina School of Medicine, Chapel Hill, North Carolina, USA

<https://doi.org/10.1289/EHP11281>

Introduction

Growing epidemiological evidence suggests that ambient air pollution, including fine particulate matter [$PM_{\leq 2.5}$ μm in aerodynamic diameter ($PM_{2.5}$)], ozone (O_3), and nitrogen dioxide (NO_2), may have detrimental effects on skin and may precipitate or exacerbate eczema, particularly during the perinatal and early childhood periods, as summarized in a review by Hassoun et al.¹ This effect is potentially due to direct oxidative stress on the skin.¹ Studies among children in Europe and Asia have demonstrated that increased exposure to outdoor air pollution is associated with increased prevalence of and visits for eczema in some populations^{2,3} but not in others.⁴ In North America, an early life cohort in Toronto, Ontario, Canada, found associations between eczema and NO_2 concentrations.⁵ Although analyses of eczema and pollution aggregated by state have been performed in the United States,⁶ to our knowledge there have not been analyses with substate exposure assessment performed to understand the impact of air pollution on eczema in a national sample of children.

Methods

We used the Medicaid Analytic eXtract (MAX) from 2009–2010, which includes Medicaid eligibility, demographic information, ZIP code tabulation area (ZCTA) of residence, and utilization records. We included individuals from birth through 18 years of age who were eligible for Medicaid from their 2009 birthday through their 2010 birthday. We defined eczema diagnosis according to the *International Classification of Diseases, Ninth Revision, Clinical Modification* (ICD-9-CM code 691.8)⁷ for any visit during the study period. We assigned exposures of long-term (2009–2010) average $PM_{2.5}$ and NO_2 and warm-season (May–September) average O_3 using a set of publicly available air pollution concentration predictions from the Center for Air, Climate, and Energy Solutions (CACES).⁸ Exposure predictions were aggregated to ZCTA averages from a census tract resolution.

Because of data quality issues of incomplete utilization data, low average months of eligibility (<9 months/year), and incomplete data on self-reported race/ethnicity [categorized as Asian, Black, Hispanic, White, Grouped, Unknown; “Grouped” combined Native American/Alaskan, Hawaiian, >1 race (Hispanic), >1 race

(non-Hispanic) due to small numbers], we excluded data from Arkansas, Colorado, Idaho, Iowa, Kansas, Maine, Massachusetts, Ohio, Pennsylvania, Rhode Island, Vermont, Washington, and West Virginia. In addition, we excluded Alaska and Hawaii because of the absence of exposure predictions. Use of the data and waiver of informed consent was approved by the Johns Hopkins School of Medicine institutional review board.

We estimated odds ratios for eczema diagnosis using generalized estimating equations with logit link and allowing for clustering by ZCTA, separately by age groups. We fit a multipollutant model with $PM_{2.5}$, O_3 , and NO_2 , as well as single-pollutant models. Models were adjusted for sex, self-reported race/ethnicity (due to associations with air pollution and with eczema rates⁹), ZCTA-level household poverty rates, season of birth,⁹ state (due to differences in Medicaid procedures by state), and unmeasured spatial confounding using 15 degrees of freedom spatial splines.¹⁰

Results

Of the 16,089,050 children included, 51% were male, 34% White, 30% Hispanic, 27% Black, and 2% Asian (Table 1). The overall

Table 1. Characteristics of Medicaid extract population by eczema status.

Characteristic ^a	Overall	Eczema	No eczema
Total sample (n)	16,089,050	432,774	15,656,276
Sex [n (%)]			
Female	7,865,322 (48.9)	222,361 (51.4)	7,642,961 (48.8)
Male	8,223,728 (51.1)	210,413 (48.6)	8,013,315 (51.2)
Self-reported race/ethnicity [n (%)]			
Asian	349,974 (2.3)	14,568 (3.4)	360,406 (2.3)
Black	4,306,462 (26.8)	151,924 (35.1)	4,154,538 (26.5)
Grouped ^b	409,783 (2.5)	8,779 (2.0)	401,004 (2.6)
Hispanic	4,803,579 (29.9)	125,175 (28.9)	4,678,404 (29.9)
White	5,451,305 (33.9)	108,955 (25.2)	5,342,350 (34.1)
Unknown	742,947 (4.6)	23,373 (5.4)	719,574 (4.6)
Age (y), [n (%)]			
<1	1,310,819 (8.1)	93,689 (21.6)	1,217,130 (7.8)
1–3	3,385,990 (21.0)	149,483 (34.5)	3,236,507 (20.7)
4–6	2,916,389 (18.1)	74,960 (18.1)	2,841,429 (18.1)
7–9	2,490,241 (15.5)	47,602 (11.0)	2,442,639 (15.6)
10–12	2,205,342 (13.7)	31,832 (7.4)	2,173,510 (13.9)
13–15	1,996,687 (12.4)	20,995 (4.9)	1,975,692 (12.6)
16–18	1,783,582 (11.1)	14,213 (3.3)	1,769,369 (11.3)
ZCTA-level poverty (%), [n (%)]			
≤5	1,512,968 (9.4)	37,880 (8.8)	1,475,088 (9.4)
6–10	3,399,517 (21.1)	87,359 (20.2)	3,312,158 (21.2)
11–15	3,691,127 (22.9)	100,988 (23.3)	3,590,139 (22.9)
16–20	2,849,931 (17.7)	76,502 (17.7)	2,773,429 (17.7)
21–25	1,818,998 (11.3)	51,449 (11.9)	1,767,549 (11.3)
26–100	2,816,509 (17.5)	78,596 (18.2)	2,737,913 (17.5)

Note: Population includes individuals eligible for Medicaid from their birthday in 2009 to their birthday in 2010. ZCTA, ZIP code tabulation area.

^aNo missing values.

^bGrouped race/ethnicity is the combination of “Native American/Alaskan,” “>1 race (Hispanic),” “>1 race (non-Hispanic),” and “Hawaiian.”

Address correspondence to Joshua P. Keller, Department of Statistics, Colorado State University, 1877 Campus Delivery, Fort Collins, CO 80523 USA. Telephone: (970) 491-7460. Email: joshua.keller@colostate.edu

J.H.D., J.P.K., and N.A.R. have no conflicts of interest to declare. R.D.P. and C.A.K. receive grant funding from the National Institutes of Health. C.A.K. holds a patent on an immunotherapy delivery method, receives royalties from UpToDate, and is a member of the American Board of Allergy and Immunology.

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehpsubmissions@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Table 2. Estimated odds ratios (95% confidence intervals) between a 1-unit difference in long-term ambient air pollution and eczema incidence (first year of life) or prevalence (all other ages), in the Medicaid population, 2009–2010, from a generalized estimating equations model.

Age (y)/model	PM _{2.5} (per 1 µg/m ³)	O ₃ (per 1 ppb)	NO ₂ (per 1 ppb)
<1			
Unadjusted single-pollutant	1.052 (1.044, 1.061)	0.996 (0.993, 0.999)	1.006 (1.001, 1.010)
Adjusted single-pollutant ^a	1.049 (1.036, 1.061)	1.004 (0.996, 1.012)	1.024 (1.019, 1.028)
Adjusted multipollutant ^b	0.987 (0.970, 1.003)	1.010 (1.003, 1.019)	1.028 (1.021, 1.034)
1–3			
Unadjusted single-pollutant	1.020 (1.012, 1.028)	0.995 (0.992, 0.998)	0.995 (0.992, 0.997)
Adjusted single-pollutant ^a	1.050 (1.040, 1.060)	1.003 (0.997, 1.010)	1.020 (1.017, 1.024)
Adjusted multipollutant ^b	1.006 (0.993, 1.020)	1.010 (1.004, 1.017)	1.020 (1.014, 1.025)
4–6			
Unadjusted single-pollutant	1.034 (1.025, 1.044)	0.994 (0.991, 0.997)	1.003 (1.000, 1.006)
Adjusted single-pollutant ^a	1.071 (1.059, 1.083)	0.995 (0.987, 1.003)	1.028 (1.024, 1.032)
Adjusted multipollutant ^b	1.012 (0.994, 1.029)	1.005 (0.997, 1.013)	1.026 (1.020, 1.032)
7–9			
Unadjusted single-pollutant	1.037 (1.028, 1.046)	0.994 (0.991, 0.998)	1.008 (1.005, 1.011)
Adjusted single-pollutant ^a	1.063 (1.050, 1.077)	0.987 (0.979, 0.996)	1.028 (1.024, 1.033)
Adjusted multipollutant ^b	0.999 (0.981, 1.017)	0.999 (0.991, 1.007)	1.028 (1.022, 1.035)
10–12			
Unadjusted single-pollutant	1.038 (1.029, 1.048)	0.996 (0.992, 1.000)	1.008 (1.005, 1.011)
Adjusted single-pollutant ^a	1.074 (1.069, 1.089)	0.989 (0.980, 0.999)	1.028 (1.023, 1.033)
Adjusted multipollutant ^b	1.021 (1.001, 1.042)	1.000 (0.990, 1.010)	1.023 (1.016, 1.030)
13–15			
Unadjusted single-pollutant	1.039 (1.029, 1.049)	1.000 (0.997, 1.004)	1.008 (1.005, 1.011)
Adjusted single-pollutant ^a	1.072 (1.056, 1.089)	0.985 (0.975, 0.995)	1.028 (1.023, 1.034)
Adjusted multipollutant ^b	1.017 (0.995, 1.040)	0.997 (0.986, 1.007)	1.024 (1.016, 1.032)
16–18			
Unadjusted single-pollutant	1.028 (1.018, 1.039)	1.001 (0.997, 1.006)	1.007 (1.004, 1.010)
Adjusted single-pollutant ^a	1.038 (1.020, 1.057)	0.982 (0.970, 0.993)	1.021 (1.015, 1.027)
Adjusted multipollutant ^b	0.985 (0.960, 1.011)	0.992 (0.980, 1.005)	1.024 (1.014, 1.033)

Note: O₃, ozone; NO₂, nitrogen dioxide; PM_{2.5}, fine particulate matter; ZCTA, ZIP code tabulation area.

^aAdjusted single-pollutant models included sex (male, female), self-reported race/ethnicity [Asian, Black, Hispanic, White, Grouped, Unknown; “Grouped” combined Native American/Alaskan, Hawaiian, >1 race (Hispanic), >1 race (non-Hispanic)], ZCTA-level household poverty rates (≤5%, 6–10%, 11–15%, 16–20%, 21–25%, 25–100%), season of birth (December–February, March–May, June–August, September–November), state, and unmeasured spatial confounding using 15 degrees of freedom spatial splines.

^bAdjusted multipollutant models included all variables from single-pollutant models and PM_{2.5}, O₃, and NO₂.

prevalence of eczema was 2.7%: highest among those in the first year of life (7.1%) and decreasing with age to 0.8% among those 16–18 years. During the study period, the median [minimum, quartile 1 (Q1), Q3, maximum] of ZCTA-average pollution levels were 8.6 µg/m³ (2.4, 7.0, 9.8, 16.1) for PM_{2.5}, 45.1 ppb (26.2, 41.8, 47.3, 58.4) for O₃, and 4.1 ppb (0.6, 3.1, 5.9, 31.2) for NO₂.

Odds of incident eczema during the first year of life were 2.8% higher (95% confidence interval: 1.021, 1.034) for each 1-ppb higher long-term NO₂ concentration, when adjusting for PM_{2.5} and O₃ (Table 2). The associations between eczema prevalence and NO₂ were similar in older age groups. There was some evidence of an association between eczema and O₃ in the youngest age groups, but this did not hold across all ages. In single-pollutant models, PM_{2.5} was associated with eczema diagnosis, but this did not persist when adjusting for NO₂ and O₃.

Discussion

In a large, geographically and racially diverse population of children, we demonstrated that long-term ambient NO₂ concentrations were associated with eczema incidence in the first year of life. Eczema, particularly in early life, is known to be associated with increased risk for multiple atopic diseases, including food allergy, asthma, and allergic rhinitis. Thus, exposures that increase eczema risk may also increase risk for other allergic diseases, and exposure reduction may decrease development of multiple allergic diseases.

Limitations of this study include the use of data that was not collected primarily for research purposes. Other uncaptured factors such as health-care seeking behaviors and secondhand smoke exposure could confound the analysis. Ambient pollution exposures were assessed as an area (i.e., ZCTA) average, and thus do not perfectly reflect personal exposures. Future work should also investigate the effects of measurement error on the estimated

associations.¹¹ For newborns, our analysis estimated associations with cumulative incidence of eczema, but for older age groups this approach is limited to associations with prevalent eczema owing to a lack of individual health-care history information prior to the study period.

In sum, in a very large U.S. cohort, we found that average exposure to NO₂ was associated with eczema prevalence. Because eczema is so central to the development of other allergic diseases, reducing exposure to NO₂ may not only prevent development of eczema but also other allergic diseases. Future research is needed to understand longitudinal associations and their mechanisms.

Acknowledgments

This work was supported by funds from the Pearl M. Stetler Research Fellowship Award, National Institutes of Health/National Institute of Allergy and Infectious Diseases (T32 AI007007 and U01AI125290 to C.A.K.) and the Johns Hopkins Innovation Grant (to C.A.K.).

References

- Hassoun Y, James C, Bernstein DI. 2019. The effects of air pollution on the development of atopic disease. *Clin Rev Allergy Immunol* 57(3):403–414, PMID: 30806950, <https://doi.org/10.1007/s12016-019-08730-3>.
- Lee JY, Lamichhane DK, Lee M, Ye S, Kwon JH, Park MS, et al. 2018. Preventive effect of residential green space on infantile atopic dermatitis associated with prenatal air pollution exposure. *Int J Environ Res Public Health* 15(1):102, PMID: 29315266, <https://doi.org/10.3390/ijerph15010102>.
- Morgenstern V, Zutavern A, Cyrys J, Brockow I, Koletzko S, Krämer U, et al. 2008. Atopic diseases, allergic sensitization, and exposure to traffic-related air pollution in children. *Am J Respir Crit Care Med* 177(12):1331–1337, PMID: 18337595, <https://doi.org/10.1164/rccm.200701-036OC>.
- Fuertes E, Sunyer J, Gehring U, Porta D, Forastiere F, Cesaroni G, et al. 2020. Associations between air pollution and pediatric eczema, rhinoconjunctivitis

- and asthma: a meta-analysis of European birth cohorts. *Environ Int* 136:105474, PMID: [31962272](#), <https://doi.org/10.1016/j.envint.2020.105474>.
5. To T, Zhu J, Stieb D, Gray N, Fong I, Pinault L, et al. 2020. Early life exposure to air pollution and incidence of childhood asthma, allergic rhinitis and eczema. *Eur Respir J* 55(2):1900913, PMID: [31806712](#), <https://doi.org/10.1183/13993003.00913-2019>.
 6. Kathuria P, Silverberg JI. 2016. Association of pollution and climate with atopic eczema in US children. *Pediatr Allergy Immunol* 27(5):478–485, PMID: [26842875](#), <https://doi.org/10.1111/pai.12543>.
 7. CDC (Centers for Disease Control and Prevention). 2013. International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM). <http://www.cdc.gov/nchs/icd/icd9cm.htm> [accessed 1 April 2018].
 8. Kim SY, Bechle M, Hankey S, Sheppard L, Szpiro AA, Marshall JD. 2020. Concentrations of criteria pollutants in the contiguous U.S., 1979–2015: role of prediction model parsimony in integrated empirical geographic regression. *PLoS One* 15(2):e0228535, PMID: [32069301](#), <https://doi.org/10.1371/journal.pone.0228535>.
 9. Dunlop JH, Keller JP, Peng RD, Keet CA. 2020. The effect of season of birth on atopic dermatitis and food allergy. *Ann Allergy Asthma Immunol* 125(2):221–223.e2, PMID: [32389781](#), <https://doi.org/10.1016/j.anai.2020.04.034>.
 10. Keller JP, Szpiro AA. 2020. Selecting a scale for spatial confounding adjustment. *J R Stat Soc Ser A Stat Soc* 183(3):1121–1143, PMID: [33132544](#), <https://doi.org/10.1111/rssa.12556>.
 11. Keller JP, Peng RD. 2019. Error in estimating area-level air pollution exposures for epidemiology. *Environmetrics* 30(8):e2573, <https://doi.org/10.1002/env.2573>.